

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 December 2000 (14.12.2000)

PCT

(10) International Publication Number
WO 00/74469 A1

- (51) International Patent Classification⁷: A01H 5/10, A23D 9/00
- (21) International Application Number: PCT/EP00/05149
- (22) International Filing Date: 5 June 2000 (05.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 09/326,500 4 June 1999 (04.06.1999) US
- (71) Applicant (for all designated States except US): **CONSEJO SUPERIOR DE INVESTIGACIONES CIENTIFICAS** [ES/ES]; Instituto de la Grasa, Avenida Padre Garcia Tejero, 4, E-41012 Sevilla (ES).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MARTINEZ-FORCE, Enrique** [ES/ES]; Lamarque de Novoa 9, 3^oA, E-41008 Sevilla (ES). **PEREZ-VICH, Begonia** [ES/ES]; Calle Ciudad Aljarafe 24, 3a-2, E-41926 Mairena del Aljarafe (ES). **FERNANDEZ-MARTINEZ, José María** [ES/ES]; Calle Teruel 10, 3^o-3, E-14012 Córdoba (ES). **GARCÉS, Rafael** [ES/ES]; Calle Dinamarca, 9, E-41907 Valencia de la Concepción (ES).
- (74) Agents: **VAN SOMEREN, Petronella, Francisca et al.**; Arnold & Siedsma, Sweelinckplein 1, NL-2517 GK The Hague (NL).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- With international search report.
 - Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 00/74469 A1

(54) Title: OIL FROM SEEDS WITH A MODIFIED FATTY ACID COMPOSITION

(57) Abstract: The invention relates to plant seeds that contain an oil having an oleic acid content of more than 5 % and less than 65 % by weight based upon the total fatty acid content, a linoleic acid content of more than 1 % and less than 65 % by weight based upon the total fatty acid content, a palmitic acid content of more than 20 % and less than 40 % by weight based upon the total fatty acid content, a stearic acid content of more than 3 % and less than 15 % based upon the total fatty acid content, and wherein the palmitoleic acid content is less than 4 % based upon the total fatty acid content, and the asclepic acid content is less than 4 % based upon the total fatty acid content. The invention further relates to the oil that can be obtained from the seeds, the use of that oil and to plants grown from the seeds or producing the seeds.

OIL FROM SEEDS WITH A MODIFIED FATTY ACID COMPOSITION**OBJECT OF THE INVENTION**

The present invention relates to oil from seeds
5 having a palmitic acid content of more than 20% and less
than 40% by weight based upon the total fatty acid
content, while the palmitoleic acid content is less than
4% based upon the total fatty acid content and the
ascleplic (n-7 isomer of oleic acid) acid content is less
10 than 4%.

In particular, the invention relates to
sunflower oil comprising an oleic acid content of more
than 5% and less than 65% by weight based upon the total
fatty acid content, a linoleic acid content less than 65%
15 by weight based upon the total fatty acid content, a
palmitic acid content of more than 20% and less than 40%
by weight based upon the total fatty acid content, a
stearic acid content of more than 3% and less than 15%
based upon the total fatty acid content, while the
20 palmitoleic acid content is less than 4% upon the total
fatty acid content and the ascleplic acid content is less
than 4% based upon the total fatty acid content.

The oil of the invention may be used for
applications in the food industry which require high
25 thermostability or plastic fats.

BACKGROUND OF THE INVENTION

Sunflower is generally cultivated for obtaining
oil which has saturated fatty acids (palmitic and
30 stearic) and unsaturated fatty acids (oleic and
linoleic), the stearic acid content is always less than
10% (Gunstone, F.D. et al. "The lipid handbook"; Chapman
and Hall, 1986), normally comprised between 3% and 7%. In
relation with the unsaturated fatty acids there are two
35 different kinds of sunflower seeds: the normal sunflower
which has a linoleic acid content between 50% and 70%
(Knowles, P.F. "Recent advances in oil crops breeding";
AOCS Proceedings, 1988) and the high oleic sunflower

which has 2-10% of linoleic acid and 75-90% of oleic acid (Soldatov, K.I. "Chemical mutagenesis in sunflower breeding"; Int. Proc. 7th Intern. Sunflower Conference, 352-357, 1976). Another high oleic sunflower line has 5 been referred by Fick (US-B1-4,627,192), with an oleic acid content of approximately 80% or greater.

Referring to saturated fatty acids, high stearic sunflower lines are disclosed in WO 95/20313. Further references to sunflower lines with high palmitic 10 acid content are WO 96/39804 and Retske et al. "Triacylglycerol composition and structure in genetically modified sunflower and soybean oils"; JAOCS 74, 989-998 (1997), European Patent Appl. 98201871.5 and Nikolova et al. "Gametocidal effect of gibberellic acid (GA₃) on 15 biochemical characteristics of sunflower seeds", Helia 15, Nr. 17, 45-50, (1992). In all these lines, the increase in palmitic acid implies higher palmitoleic acid content, always over 4%, and where the two "cis" isomers n-7 octadecenoic acid (asclepic) and n-9 octadecenoic 20 (oleic) acid have been analyzed, an increase in the n-7 (asclepic acid) isomer has been observed.

Table 1 shows the fatty acid composition for all this indicated sunflower oil varieties.

25 Table 1

Reference	Line	Fatty acids composition (%)								
		16:0	16:1	16:2	18:0	18:1	18:1A	18:2	20:0	22:0
Gunstone	Normal	6	-	-	5	18	-	68	-	1
	Oleic	5	-	-	4	88	-	2	-	1
Nikolova,1992		29	5	*	2	9	*	55	*	*
	Reske, 1997	25	6	*	3	60	*	4	-	2
		27	4	*	3	17	*	47	-	1
CAS-5		31	5	1	3	5	6	47	-	1
	CAS-12	30	7	-	2	52	4	3	-	2

* These fatty acids were not determined in those papers.

- = Traces.

The saturated fatty acid content of an oil is directly related with the physical and chemical characteristics thereof. In case that said content is sufficiently high, the oil can be a solid at room temperature like some animal fats. Normal sunflower oil is always a liquid under said conditions. In the food industry like for the production of confectionery or margarine, animal fats or hydrogenated vegetable fats are usually used because a solid or semisolid product is required. By means of hydrogenation unsaturated fatty acids are converted into saturated fatty acids. Animal fats as well as hydrogenated fats are not very recommendable from a nutritional point of view (Chow, C.K. "Fatty acids in food and their health implications", Dekker, N.Y., 1992). Animal fats have a relatively high cholesterol content. Too much cholesterol in the diet may be detrimental to the health. Therefore animal fats have been substituted in the last years by hydrogenated vegetable fats which do not contain cholesterol.

However, said hydrogenated fats present another problem derived from the hydrogenation process. In said process positional isomerization (shift of double bonds) and stereo-chemical transformations (formation of "trans" isomers) take place. Isomers are produced in an amount of up to 30%-50% of the total fatty acids amount. These isomers are not very healthy from a nutritional point of view (Wood, R. "Biological effects of geometrical and positional isomers of monounsaturated fatty acids in humans", Dekker, N.Y. 1990; Willet, W.C. & Ascherio, A., "Trans Fatty Acids: Are the effects only marginal?", American Journal of Public Health, Vol. 84, 5, 1994). Therefore, the use of hydrogenated fats in the food industry should be avoided.

As previously referred, the increase in palmitic acid implies higher palmitoleic acid contents, always over 4% (see WO 96/39804). These oils are useful for food industry which requires high thermostability, but the presence of the indicated palmitoleic acid contents is

still undesirable. Studies carried out on macadamia oil, which has 4% palmitoleic acid content indicate a negative effect on plasmatic cholesterol when compared with palmitic and oleic (Nestel et al., "Effects of increasing dietary palmitoleic acid compared with palmitic and oleic acids on plasma lipids of hypercholesterolemic men", Journal Lipid Research, vol. 35, pp. 656-662, 1994). This oil has also higher asclepic acid contents (n-7 isomer of octadecanoic acid) than other normal sunflower oil that have 0.6% or other vegetable oils, like soybean or rape which have 0.8 and 0.9% respectively (Mukherjee K.D. and Kiewitt I., "Formation of (n-9) and (n-7) cis-monounsaturated fatty acids in seeds of higher plants", Planta, vol. 149, pp. 461-463.

It can be concluded that an oil having higher palmitic and stearic acid contents than normal sunflower oil, but maintaining reduced levels of palmitoleic and asclepic acids would meet all the requirements for food industry implying high thermostability and plasticity to be spread.

BRIEF DESCRIPTION OF THE INVENTION

The present invention relates to oil from seeds comprising an oleic acid content of more than 5% and less than 65% by weight based upon the total fatty acid content, a linoleic acid content of more than 1% and less than 65% by weight based upon the total fatty acid content, a palmitic acid content of more than 20% and less than 40% by weight based upon the total fatty acid content, a stearic acid content of more than 3% and less than 15% based upon the total fatty acid content, while the palmitoleic acid content is less than 4% based upon the total fatty acid content and the asclepic acid content is less than 4% based upon the total fatty acid content. Preferably, the palmitoleic acid content is less than 3% based upon the total fatty acid content and the asclepic acid content is less than 3% based upon the total fatty acid content.

The oil from seeds according to the present invention has an oleic acid content which is at least 40% by weight based upon the total fatty acid content and a linoleic acid content which is less than 18%. The total level of saturated fatty acids in the oil is at least 26%, preferably higher than 35%, most preferably higher than 45% based upon the total fatty acids content.

In reference to the position of the fatty acid groups in the triacylglycerols (TAG), the oil of the invention has less than 10%, preferably less than 5% by weight of the saturated fatty acid groups in the 2 position of the TAGs.

The invention in particular relates to sunflower oil, which is extracted from sunflower seeds obtained by crossing sunflower seeds of the mutant sunflower line IG-1297M deposited on 20 January 1998 with ATCC under deposit accession number ATCC-209591 with the mutant sunflower line CAS-3, deposited on 14 December 1994 with the ATCC under deposit accession number ATCC-75968.

The invention further relates to sunflower seeds comprising a sunflower oil with a fatty acid composition as referred herein above and to a method for preparing sunflower seeds, comprising the steps of:

a) crossing sunflower seeds of the mutant sunflower line IG-1297M deposited on 20 January 1998 with ATCC under deposit accession number ATCC-209591 with the mutant sunflower line CAS-3, deposited on 14 December 1994 with the ATCC under deposit accession number ATCC-75968;

b) self-pollinating F1 progeny plants of step a) for at least two generations to produce inbred plants;

c) selecting from the progeny of step b) plants with seeds containing an oil having a palmitic acid higher than 20%, palmitoleic acid content of less than 4% and an asclepic acid content of less than 3%;

d) collecting progeny seeds from step c) and optionally;

e) repeating the cycle of culturing, selection and collection of seeds.

The invention further relates to plants producing the seeds of the invention and to plants
5 obtainable by growing the seeds of the invention and their progeny.

The sunflower oil, prepared by extracting said sunflower seeds may be used in roasting, cooking, frying, baking and in general at high temperature conditions
10 which constitute heating by any means at temperatures of at least 70°C. Said oil may also be used in the production of edible fats or fat mixtures, such as margarine, shortening or vegetable-dairy as well as in confectionery and bakery.

15

DETAILED DESCRIPTION OF THE INVENTION

The method for preparing seeds with a modified fatty acid composition comprises mutagenesis of seeds with a suitable mutagenic agent. The mutagenesis will
20 produce inheritable genetic changes in the DNA of the seeds. According to the invention it was possible after several different experiments to select some treatments that produce a high number of genetic modifications in the genes that control the seed fatty acid biosynthesis.
25 These treatments comprise the use of sodium azide or an alkylating agent, like ethyl methane sulfonate. Of course any other mutagenic agent having the same or similar effects may also be used.

Then, the next seed generation was analyzed with
30 a new methodology described in Garcés, R. and Mancha, M. "One step lipid extraction and fatty acid methyl esters preparation from fresh plant tissues"; Analytical Biochemistry, 211:139-143, 1993. This allowed for the detection of seeds with modifications in the composition
35 of any fatty acid. Selected seeds showing a desirable fatty acid composition have been cultivated to the fifth generation showing that this new genetic trait is inheritable and stable and independent of growth

conditions. In the method of the invention the parent seeds are for example treated during 2 hours at room temperature with a solution of 70 mM ethyl methane sulfonate in water, or during 2 hours at room temperature 5 with a solution of 2 mM sodium azide in water. Further, the mutation and selection steps may be followed by conventional plant improvement techniques thus leading to seeds having a desirable fatty acids content.

The seeds of the invention may be subjected to 10 one or more further mutation treatments. Another way of obtaining mutagenized seeds consists of submitting the seeds to X-rays action, growing the treated seeds, self-pollination and further analysis of the fatty acid content. Further growing and selection steps will lead to 15 plants with the desired new character fixed.

Sunflower oil having the desirable fatty acid composition may be prepared by extraction from sunflower seeds of the invention in any manner known to the person skilled in the art. Such extraction methods are well 20 known and for example described in "Bailey's industrial oil and fat products", Vol.2, Chapter 3; 4th Edition, John Wiley and Sons, New York (1982).

By the referred methods seeds and oil having high stearic acid and high palmitic acid content can be 25 obtained. High palmitic acid content normally implies high palmitoleic acid content which is not desirable from a nutritional point of view, as previously indicated. However, biochemical research on sunflower mutant lines indicates that the high stearic mutant has less stearyl 30 desaturase activity over palmitoyl-ACP than other sunflower mutant lines. Crossing a high stearic line (CAS-3) with a high palmitic line (CAS-12) and selecting in search of different fatty acid compositions, it turned out that in certain F2 generations that amounts of 35 palmitoleic and asclepic acid decreased. Thus, the desaturation of palmitic into palmitoleic in the high palmitic acid mutants could be reduced introducing the

stearoyl desaturase enzymatic activity of the high stearic mutant lines.

The invention is further illustrated by means of the following examples.

5

EXAMPLE 1

Sunflower seeds RDF-1-532 (Sunflower Collection of Instituto de Agricultura Sostenible, CSIC, Córdoba, Spain), which have 4% to 7% stearic acid content were
10 mutagenized with a solution of 70 mM of ethyl methane sulfonate (EMS) in water. The treatment was performed at room temperature during 2 hours while shaking (60 rpm). After mutagenesis the EMS solution was discarded and seeds were washed during 16 hours under tap water.

15 Treated seeds were germinated in the field and plants were self-pollinated. The seeds collected from these plants were used to select new sunflower lines with modifications in the fatty acid composition. By using the method of Garcés, R. and Mancha, M, referred to herein
20 above, the seed fatty acid composition was determined by gas-liquid chromatography, after converting the fatty acids into their corresponding methyl esters.

A first plant with 9 to 17% stearic acid content in the oil was selected. The progeny was cultivated for
25 five generations wherein the stearic acid content increased and the new genetic trait became stably fixed in the genetic material of the seed. This line is called CAS-3. A selected sample of this line was analyzed resulting in a stearic acid content of 26% (Table 2). The
30 minimum and the maximum stearic acid content of the line were 19 and 35%, respectively. The stearic acid content of oil extracted from seeds from this cell line may thus lie between 19 and 35%.

35 **EXAMPLE 2**

Sunflower seeds RDF-1-532 were mutagenized with sodium azide, at a concentration of 2 mM in water. The treatment was performed at room temperature during two

hours while shaking. Then the mutagenesis solution was discarded and seeds were washed during 16 hours with tap water.

Seeds were planted in the field and plants were self-pollinated. Seeds from these plants were collected, and the fatty acid composition was determined by gas-liquid chromatography, after converting the fatty acids into their corresponding methyl esters using the method described in Example 1. Seeds from a plant having around 10% stearic acid in the oil were selected and cultivated for five generations. During this procedure the stearic acid content was increased and the new genetic trait fixed. This line is called CAS-4. A selected sample of this line was analyzed resulting in a stearic acid content of 16.1%. The minimum and the maximum values were 12 and 19% respectively.

EXAMPLE 3

5000 dry sunflower seeds were mutagenized by treatment with X-rays 300 cGy/min, beam 200 kV, 18 mA⁻¹ and dose 160 Gy with a Siemens Stabilipan (Erlangen, Germany), seeds were grown in spring in the field. Self-pollinated plants were collected individually and seeds analyzed for their fatty acid content. Seeds with at least three times more saturated fatty acid content than the standard deviation for the specific fatty acid were selected and successively grown until the new character was fixed. Several putative new mutant lines were selected by this method. After further selection for triacylglycerol composition line IG-1297M was selected.

EXAMPLE 4

Sunflower plants were grown from the sunflower seeds of CAS-3 according to Example 1. Sunflower plants were grown from the sunflower seeds of IG-1297M according to Example 3.

The lines were crossed. The plants were assisted by artificial pollination in order to ensure adequate

seed production occurred. The F1 was produced on the IG-1297M and harvested. F2 IG-1297M parent seeds with a high oleic acid background having more than 20% palmitic acid and less than 4% of both palmitoleic and asclepic acid
5 were selected.

Although the oil produced by these selected lines is the oil of the present invention, the level of production is limited, therefore fixed inbred lines evidencing seeds with these oil profiles are desirable.
10 These homozygous fixed high oleic, high palmitic, low palmitoleic, low asclepic inbred lines can then be crossed to form hybrid seed, which will produce F2 seed evidencing the desired oil traits of the present invention.

15 Toward this end the F1 seeds were planted and produced plants were selfed in isolated conditions and F2 seed was produced. The F2 seed, called QQ-3598-M was tested for the four traits: high palmitic, high oleic, low palmitoleic, and low asclepic. The remaining portion
20 of the seeds evidencing these traits was employed to grow plants to form F3 seed. The selfing and screening and selection process is repeated to develop the fixed homozygous QQ-3598-M line, having the following fatty acid profile: C16:0 30,5%; C18:0 9,6%; C18:1 47,2%; C18:2
25 6,7%; C16:1 2,1%; C18:1A 1,1%, and less than 1% of other minor fatty acids.

Once the trait is fixed, similar QQ-3598-M lines can cross to form hybrid seed having the desired traits. This characteristic fatty acid profile is an inheritable
30 trait and is fairly independent from the growing conditions.

CLAIMS

1. Plant seeds that contain an oil having an oleic acid content of more than 5% and less than 65% by weight based upon the total fatty acid content, a linoleic acid content of more than 1% and less than 65% by weight based upon the total fatty acid content, a palmitic acid content of more than 20% and less than 40% by weight based upon the total fatty acid content, a stearic acid content of more than 3% and less than 15% based upon the total fatty acid content, characterized in that

- the palmitoleic acid content is less than 4% based upon the total fatty acid content; and
- the asclepic acid content is less than 4% based upon the total fatty acid content.

2. Seeds according to claim 1, characterized in that the palmitoleic acid content in the oil is less than 3% based upon the total fatty acid content.

3. Seeds according to claim 1 or 2, characterized in that the asclepic acid content in the oil is less than 2% based upon the total fatty acid content.

4. Seeds according to claims 1-3, characterized in that the oleic acid content in the oil is at least 40% by weight based upon the total fatty acid content.

5. Seeds according to claims 1-4, characterized in that the total level of saturated fatty acids in the oil is at least 24% by weight based upon the total fatty acid content.

6. Seeds according to claim 5, characterized in that the total level of saturated fatty acids in the oil is at least 35% by weight based upon the total fatty acids content.

7. Seeds according to claim 6, characterized in that the total level of saturated fatty acids in the oil is at least 45% by weight based upon the total fatty acids content.

8. Seeds according to claims 1-7, characterized in that the linoleic acid content in the oil is less than 18% by weight based upon the total fatty acids content.

9. Seeds according to claims 1-8, characterized in that the oil has less than 10% by weight of the saturated fatty acid groups in the 2 position of the triacylglycerol molecules of the oil.

10. Seeds according to claim 9, characterized in that the oil has a maximum of 5% of the saturated fatty acid groups in the 2 position of the triacylglycerol molecules of the oil.

11. Seeds according to claims 1-10, which seeds are sunflower seeds.

12. Seeds according to claim 11, obtainable by crossing sunflower seeds of the mutant sunflower line IG-1297M deposited on 20 January 1998 with ATCC under deposit accession number ATCC-209591 with the mutant sunflower line CAS-3, deposited on 14 December 1994 with the ATCC under deposit accession number ATCC-75968.

13. Oil having an oleic acid content of more than 5% and less than 65% by weight based upon the total fatty acid content, a linoleic acid content of more than 1% and less than 65% by weight based upon the total fatty acid content, a palmitic acid content of more than 20% and less than 40% by weight based upon the total fatty acid content, a stearic acid content of more than 3% and less than 15% based upon the total fatty acid content, characterized in that the palmitoleic acid content is less than 4% based upon the total fatty acid content, and; and the asclepic acid content is less than 4% based upon the total fatty acid content.

14. Oil according to claim 13, obtainable by extracting seeds as claimed in claims 1-12.

15. Oil according to claims 13 and 14, which oil is a sunflower oil.

16. Plants producing the seeds as claimed in claims 1-12.

17. Plants, obtainable by growing seeds as claimed in claims 1-12.

18. Plants according to claims 16 and 17, which plants are sunflower plants.

5 19. Progeny of the plants according to claims 16-18.

20. Method for preparing sunflower seeds as claimed in 12, comprising the steps of:

a) crossing sunflower seeds of the mutant
10 sunflower line IG-1297M deposited on 20 January 1998 with ATCC under deposit accession number ATCC-209591 with the mutant sunflower line CAS-3, deposited on 14 December 1994 with the ATCC under deposit accession number ATCC-75968;

15 b) self-pollinating F1 progeny plants of step a) for at least two generations to produce inbred plants.

c) selecting from the progeny of step b) plants with seeds containing an oil having a palmitic acid content of more than 20%, a palmitoleic acid content of
20 less than 4% and an asclepic acid content of less than 3%.

d) collecting progeny seeds from step c) and optionally

e) repeating the cycle of self-pollination,
25 selection and collection of seeds.

21. Method for preparing an oil according to claim 13-15, by extracting seeds as claimed in claims 1-12.

22. Use of oil according to claims 13-15 at high
30 temperature conditions.

23. Use of oil as claimed in claim 22, wherein the high temperature conditions constitute baking.

24. Use of oil as claimed in claim 22, wherein the high temperature conditions constitute cooking.

35 25. Use of oil as claimed in claim 22, wherein the high temperature conditions constitute roasting.

26. Use of oil as claimed in claim 22, wherein the high temperature conditions constitute heating by any means at temperatures of at least 70°C.

27. Use of the oil according to claims 13-15 in 5 the production of edible fats or fat mixtures, such as margarine or vegetable-dairy.

28. Use of the oil according to claims 13-15 in confectionery or bakery.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/05149

A. CLASSIFICATION OF SUBJECT MATTER
 IPC 7 A01H5/10 A23D9/00

According to international Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A01H A23D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, FSTA

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	AGGARWAL M ET AL: "The effect of Cd-2+ on lipid components of sunflower (Helianthus annuus L.) seeds." PLANT FOODS FOR HUMAN NUTRITION (DORDRECHT), vol. 47, no. 2, 1995, pages 149-155, XP000952476 ISSN: 0921-9668 tables 2-4 ---	1,13,16, 20,22-28
A	OSORIO J ET AL: "MUTANT SUNFLOWERS WITH HIGH CONCENTRATION OF SATURATED FATTY ACIDS IN THE OIL" CROP SCIENCE,US,CROP SCIENCE SOCIETY OF AMERICA, MADISON, WI, vol. 35, 1995, pages 739-742, XP000677842 ISSN: 0011-183X table 1 ---	1,13,16, 20,22-28
-/--		

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

13 October 2000

Date of mailing of the international search report

30/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Fonts Cavestany, A

INTERNATIONAL SEARCH REPORT

Intern 1al Application No

PCT/EP 00/05149

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; February 1999 (1999-02) MARTINEZ-FORCE ENRIQUE ET AL: "Enzymatic characterisation of high-palmitic acid sunflower (Helianthus annuus L.) mutants." Database accession no. PREV199900194881 XP002150054 abstract & PLANTA (BERLIN), vol. 207, no. 4, February 1999 (1999-02), pages 533-538, ISSN: 0032-0935</p> <p style="text-align: center;">---</p>	1, 13, 16, 20, 22-28
A	<p>DATABASE BIOSIS 'Online! BIOSCIENCES INFORMATION SERVICE, PHILADELPHIA, PA, US; 1997 RESKE JOHANNA ET AL: "Triacylglycerol composition and structure in genetically modified sunflower and soybean oils." Database accession no. PREV199799705973 XP002150055 cited in the application abstract & JOURNAL OF THE AMERICAN OIL CHEMISTS' SOCIETY, vol. 74, no. 8, 1997, pages 989-998, ISSN: 0003-021X</p> <p style="text-align: center;">---</p>	1
A	<p>US 5 710 366 A (COLE GLENN S ET AL) 20 January 1998 (1998-01-20) cited in the application claims; tables</p> <p style="text-align: center;">---</p>	1
A	<p>WO 95 20313 A (OSORIO JORGE ;GARCES RAFAEL (ES); MANCHA MANUEL (ES); CONSEJO SUPE) 3 August 1995 (1995-08-03) cited in the application claims; tables</p> <p style="text-align: center;">---</p>	1
A	<p>DATABASE FSTA 'Online! INTERNATIONAL FOOD INFORMATION SERVICE (IFIS), FRANFURT/MAIN, DE; FERNANDEZ-MARTINEZ J M ET AL: "Sunflower mutant containing high levels of palmitic acid in high oleic background." Database accession no. 1998-03-n0113 XP002150056 abstract & EUPHYTICA 1997 INST. DE AGRIC. SOSTENIBLE, CSIC, APDO. 4084, 14080 CORDOBA, SPAIN, vol. 97, no. 1, pages 113-116,</p> <p style="text-align: center;">-----</p>	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/05149

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 5710366 A	20-01-1998	AU 705124 B	13-05-1999
		AU 6261096 A	30-12-1996
		CA 2221639 A	19-12-1996
		EP 0836376 A	22-04-1998
		HU 9900889 A	28-07-1999
		JP 11509405 T	24-08-1999
		SK 166897 A	06-05-1998
		WO 9639804 A	19-12-1996
		US 5872271 A	16-02-1999

WO 9520313 A	03-08-1995	ES 2082716 A	16-03-1996
		ES 2082717 A	16-03-1996
		AU 706751 B	24-06-1999
		AU 1705495 A	15-08-1995
		BG 62751 B	31-07-2000
		BG 100734 A	29-08-1997
		CA 2181261 A	03-08-1995
		DE 69511844 D	07-10-1999
		DE 69511844 T	16-12-1999
		EP 0741511 A	13-11-1996
		EP 0934692 A	11-08-1999
		ES 2095799 A	16-02-1997
		HU 75362 A	28-05-1997
		ES 2095800 A	16-02-1997
